arXiv:1005.5344v1 [math.NA] 28 May 2010

Numerical convergence of a one step approximation of an intrgro-differential equation

Samir Kumar Bhowmik *
KdV Institute for Mathematics, University of Amsterdam
Amsterdam, Netherlands, S.K.Bhowmik@uva.nl
and
Department of Mathematics, University of Dhaka
Dhaka 1000, Bangladesh, bhowmiksk@gmail.com.

May 31, 2010

Abstract

We consider a linear partial integro-differential equation that arises in the modeling of various physical and biological processes. We study the problem in a spatial periodic domain. We analyze numerical stability and numerical convergence of a one step approximation of the problem with smooth and non-smooth initial functions.

Keywords: stability; convergence; smoothness; full discrete problem; infinite domain; periodic domain.

1 Introduction

Study in convolution model of phase transitions (initial value problem) is of ongoing interest. Many scientific problems have been modeled by reaction diffusion equation and advection reaction diffusion equations. A lot of

---

[*] The Author would like to thank Professor Dugald B. Duncan for his valuable advise and kind help.

models consider nonlocal type diffusion operators [2, 3, 4, 5, 10, 13]. In this article we analyze stability, accuracy and rate of convergence of a simple approximation of such a linear partial integro-differential equation (IDE)

$$u_t(x,t) = \varepsilon \int_\Omega J(x-y)\left(u(y,t)-u(x,t)\right)dy = \mathbb{L}u, \tag{1}$$

which is the linear part of the IDE [2, 3, 6, 7, 9, 10, 11, 12]

$$u_t = \varepsilon \left(\int_\Omega J(x-y)u(y,t)dy - u(x,t)\int_\Omega J(x-y)dy \right) + f(u). \tag{2}$$

Here the initial condition $u(x,0) = u_0(x), x \in \Omega$ where $\Omega \subseteq \mathbb{R}$, $f(u)$ is a bistable nonlinearity for the associated ordinary differential equation $u_t = f(u)$ and $J(x-y)$ is a kernel that measures interaction between particles at positions x and at y. Here it is assumed that the effect of close neighbours x and y is greater than that from more distant ones; the spatial variation is incorporated in J. We assume that J is a non-negative function satisfies smoothness, symmetry and decay conditions.

We present the stability and convergence of the rescaled integro-differential equation model in a periodic domain. Here we consider spacial $[0,1]$ periodic domain. For simplicity of notations, from here J, u mean functions in the periodic domain and J^∞, u^∞ mean functions defined in the infinite domain. If we choose spatially one-periodic initial data $u(x,0)$, then for all $x \in \mathbb{R}$ and $t \in \mathbb{R}_+$

$$u(x,t) = u(x+1,t).$$

Then a with kernel function $J^\infty(x)$, (1) can be written as

$$\begin{aligned} u_t &= \varepsilon \int_{\mathbb{R}} J^\infty(x-y)\left(u(y,t)-u(x,t)\right)dy \\ &= \varepsilon \int_0^1 J(x-z)\left(u(z,t)-u(x,t)\right)dz, \end{aligned} \tag{3}$$

where

$$J(x) = \sum_{r=-\infty}^{\infty} J^\infty(x-r) \tag{4}$$

and $x \in [0,1]$. Two sample kernels are shown in Figure 1. Here we consider $\varepsilon = 1$ only since it does not affect our analysis.

In [5], the author study stability and accuracy of an one step approximation the IDE considering infinite spatial domain. They first write a discrete



Figure 1: Examples of kernel functions in $[0, 1]$ periodic domain defind by (4) where (a) $J^{\infty}(x) = \sqrt{\frac{100}{\pi}} e^{-100x^2}$, (b) $J^{\infty}(x) = \frac{c}{2} e^{-c|x|}$, $c = 10$.

equivalent of the integral equation, then use the forward Euler method for time discretisation. They present some numerical results to demonstrate the rate of convergence for smooth and non-smooth initial functions.

In [10], Duncan et. al. consider (2) in a spatial periodic $[0, 1]$ and $[0, 1]^2$ domains. They approximate the problem using piecewise constant basis function with collocation and mid point quadrature rule for space discretisation, then they use some standard ode solver for time integration. They present some numerical results to demonstrate their scheme. In [8], the author consider several local and nonlocal operators that contains (1). He approximate the models using finite difference schemes and discuss some stability issues.

Here we consider the problem (3) in a spatial periodic domain, and study stability and accuracy of a simple scheme using the Fourier series and the discrete Fourier transform definitions. We Organize our article in the following way. We present a simple space time discretisation in Section 2 followed by stability analysis of the scheme in Section 3. We analyze accuracy of the scheme in Section 4 considering smooth initial function, whereas in Section 5, we discuss the same considering non-smooth initial function. We study accuracy of the semi-discrete time dependent scheme in Section 6.

We finish our study in Section 7 with some numerical experiments and discussions.

2 Numerical approximation

We approximate (1) as

$$\frac{dU_j(t)}{dt} = h\sum_{k=0}^{N-1} J(x_j - x_k)(U_k - U_j) \tag{5}$$

for each $j = 0, 1, 2, \cdots, N-1$ where $U_j(t) \approx u(x_j, t)$ and $x_j = jh$. Applying Euler's method we can again approximate the semi-discrete problem (5) as

$$U_j^{n+1} - U_j^n = \Delta t h \sum_{k=0}^{N-1} J(x_j - x_k)\left(U_k^n - U_j^n\right)$$

$$U_j^{n+1} = U_j^n\left(1 - h\Delta t\sum_{r=0}^{N-1} J(x_j - x_r)\right) + h\Delta t\sum_{r=0}^{N-1} J(x_j - x_r)U_r^n. \tag{6}$$

We use Fourier series and discrete Fourier transforms throughout. Before the main discussion let us introduce some necessary definitions and theorems. The Discrete Fourier Transform for a periodic function (DFT) can be defined as

$$\tilde{U}_k = h\sum_{j=0}^{N-1} u(x_j)e^{-i2\pi k x_j}$$

where $u(x,t)$ is a periodic function with period 1 and its inverse Fourier transform is defined as

$$u_j = h\sum_{k=0}^{N-1} \tilde{U}_k e^{i2\pi k x_j} \quad \text{or} \quad u_j = h\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \tilde{U}_k e^{i2\pi k x_j}.$$

For any function periodic function $f(x)$ with period 1, its Fourier series can be defined as

$$f(x) = \sum_{n=-\infty}^{\infty} \hat{f}_n e^{i2\pi n x} \quad \text{where} \quad \hat{f}_n = \int_0^1 f(x)e^{-i2\pi n x}dx.$$

For any 1-periodic complex-valued function $u(x)$ the relation

$$\sum_{n=-\infty}^{\infty} |\hat{u}_n|^2 = \int_0^1 |u(x)|^2 dx$$

where $\hat{u}_n$ are the Fourier coefficients of $u(x)$ and for a real-valued function $u(x)$

$$\frac{\hat{u}_0^2}{4} + \frac{1}{2}\sum_{n=1}^{\infty}\left(\hat{u}_{-n}^2 + \hat{u}_n^2\right) = \int_0^1 u^2(x)dx$$

is called Parseval's relation for the Fourier series where a_n, b_n are Fourier coefficients. For the Discrete Fourier Transform of $u(x)$, the relation

$$\sum_{j=0}^{N-1}|u_j|^2 = h\sum_{k=0}^{N-1}|\tilde{U}_k|^2 \quad or \quad h\textstyle\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}}|\tilde{U}_k|^2$$

is called Parseval's relation where $\tilde{U}_k$ is the DFT of u_j, both have the same length N.

Now we will discuss the relation between the DFT and the Fourier coefficients. From the DFT definition $\tilde{U}_k = h\sum_{j=0}^{N-1} u(jh)e^{-i2\pi kx_j}$, the Fourier coefficients are

$$\hat{u}_k = \int_0^1 u(x)e^{-i2\pi kx}dx$$

and the corresponding Fourier series is

$$u(x) = \sum_{k=-\infty}^{\infty}\hat{u}_k e^{i2\pi kx}.$$

Thus

$$\begin{aligned}\tilde{U}_k &= h\sum_{j=0}^{N-1}\sum_{r=-\infty}^{\infty}\hat{u}_r e^{i2\pi rx_j}e^{-i2\pi kx_j} \\ &= h\sum_{r=-\infty}^{\infty}\hat{u}_r N\delta_N(r-k) = \sum_{m=-\infty}^{\infty}\hat{u}_{k+mN} \qquad (7)\end{aligned}$$

where $r-k = mN$ and

$$\delta_N(j) = \begin{cases} 1 & j = mN \text{ for some } m \in \mathbb{Z} \\ 0 & \text{otherwise.}\end{cases} \qquad (8)$$

The relation (7) is known as the discrete Poisson sum formula [15]. The Fourier coefficients of the kernel function in $[0,1]$ are given by

$$\hat{J}_j = \int_0^1 J(x)e^{-i2\pi jx}dx = \int_{-\infty}^{\infty} J^{\infty}(y)e^{-i2\pi jy}dy = \hat{J}^{\infty}(2\pi j) \qquad (9)$$

is the relation between the Fourier coefficient of $J(x)$ defined in (4) and the continuous Fourier transform of $J^{\infty}(x)$.

Multiplying both sides of (6) by $he^{-i2\pi kx_j}$ and summing over j

$$\begin{aligned} h\sum_{j=0}^{\mathbb{N}-1} U_j^{n+1} e^{-i2\pi kx_j} &= h\sum_{j=0}^{N-1} U_j^{n} e^{-i2\pi kx_j}\left(1 - h\Delta t\sum_{r=0}^{N-1} J(x_j - x_r)\right) \\ &\quad + h\sum_{r=0}^{N-1} e^{-i2\pi kx_j} U_r^n \left(h\Delta t\sum_{j=0}^{N-1} J(x_j - x_r) e^{-i2\pi kx_j}\right). \end{aligned}$$

That is,

$$\begin{aligned} \tilde{U}_k^{n+1} &= \tilde{U}_k^{n}\left(1 - h\Delta t\sum_{r=0}^{N-1} J(x_j - x_r) + h\Delta t\sum_{j=0}^{N-1} J(x_j - x_r) e^{-i2\pi kx_j}\right) \\ &= g(h, \Delta t, k)\tilde{U}_k^{n} \end{aligned}$$

and clearly

$$\tilde{U}_k^{n} = g^{n}(h, \Delta t, k)\tilde{U}_k^{0}, \tag{10}$$

where

$$g(h, \Delta t, k) = 1 + h\Delta t\sum_{j=0}^{N-1} J(x_j - x_r)\left(e^{-i2\pi kx_j} - 1\right) = 1 + \Delta t\left(\tilde{J}_k - \tilde{J}_0\right)$$

since J is 1-periodic.

3 Stability analysis

To show the stability of the scheme we need some reasonable restrictions on $J(x)$. We will examine properties of the DFT of $J(x)$ under some reasonable hypothesis on the function $J(x)$. We use the bounds obtained below to get a bound on $g(h, \Delta t, k)$.

Lemma 1. *Assume that*

A1. $J(x) \geq 0$.

A2. $J(x) = J(-x)$.

A3. $\int_{\Omega} J(x)dx = 1$.

A4. *$\frac{d}{dx}J(x) < 0$ for $x \in (0, \frac{1}{2})$*

A5. *$\hat{J}_k \geq 0$ for $-\frac{N}{2} + 1 \leq k \leq \frac{N}{2}$*

then $0 \leq \tilde{J}_0 - \tilde{J}_k \leq 2$, $\hat{J}_0 - \hat{J}_k \leq 2$ where $-\frac{N}{2} + 1 \leq k \leq \frac{N}{2}$. Furthermore, $\tilde{J}_k \geq 0$.

Proof. We have

$$\tilde{J}_k = h \sum_{r=0}^{N-1} J(x_r) e^{-ik2\pi x_r} \quad \text{and} \quad \tilde{J}_0 = h \sum_{r=0}^{N-1} J(x_r).$$

So

$$\tilde{J}_0 - \tilde{J}_k = h \sum_{r=0}^{N-1} J(x_r) \left(1 - e^{-ik2\pi x_r}\right).$$

For simplicity from here we take N to be even. Using the symmetry of $J(x)$ in $[0, 1]$

$$\tilde{J}_0 - \tilde{J}_k = 2h \sum_{r=0}^{\frac{N}{2}} J(x_r) \left(1 - \cos(k2\pi x_r)\right) \geq 0$$

as $h > 0$ and $J \geq 0$. Also $1 - \cos(k2\pi x_r) \leq 2$. Thus

$$\tilde{J}_0 - \tilde{J}_k = 4h \sum_{r=0}^{\frac{N}{2}} J(x_r) \leq 4 \int_0^{\frac{1}{2}} J(x)dx = 2 \int_0^1 J(x)dx = 2.$$

So we conclude $0 \leq \tilde{J}_0 - \tilde{J}_k \leq 2$. The result $\hat{J}_0 - \hat{J}_k \leq 2$ follows from the definition of J and by using similar steps to those of [4]. □

Lemma 2. *If $J(x)$ satisfies **A1** - **A3** then $1 \geq g(h, \Delta t, k) \geq 1 - C\Delta t$ for some $C > 0$ and if $J(x)$ satisfies **A4** - **A5** from Lemma 1 as well then $C = 2$ and*

$$|g(h, \Delta t, k)| \leq 1 \quad \text{for all} \quad 0 < \Delta t \leq \Delta t^* = \tfrac{2}{C}.$$

Proof. Proof of this Lemma follows directly from Lemma 1. □

The stability result follows from the following theorem.

Theorem 1. *If $J(x)$ is a periodic function in $[0, 1]$ and satisfies **A1** - **A5**, then there exists $\Delta t^* > 0$ given by Lemma 2 such that $\|U^n\|_h \leq \|U^0\|_h$ for all $0 < \Delta t \leq \Delta t^*$ and $n \geq 0$.*

Proof. We have

$$\begin{aligned}\|U^n\|_h^2 &= h\sum_{j=0}^{n-1}|U_j^n|^2 = h^2\sum_{j=0}^{N-1}\left|\tilde{U}_j^n\right|^2 = h^2\sum_{j=0}^{N-1}\left|g^n(h,\Delta t,k)\tilde{U}_j^0\right|^2 \\ &\leq h^2\sum_{j=0}^{N-1}|\tilde{U}_j^0|^2 \leq h\sum_{j=0}^{N-1}|U_j^0|^2 = \|U_j^0\|_h^2,\end{aligned}$$

which gives the result. □

4 Convergence analysis of the fully discrete approximation

Let the Fourier series of $u(x,t)$ be

$$u(x,t) = \sum_{j=-\infty}^{\infty}\hat{u}_j(t)e^{ij2\pi x} \tag{11}$$

where

$$\hat{u}_j(t) = \int_0^1 u(x,t)e^{-ij2\pi x}dx$$

and let the Fourier series expansion of $J(x)$ be

$$J(x) = \sum_{j=-\infty}^{\infty}\hat{J}_j e^{ij2\pi x} \tag{12}$$

where

$$\hat{J}_j = \int_0^1 J(x)e^{-ij2\pi x}dx.$$

Substituting (11)-(12) in (1)

$$\begin{aligned}\sum_{j=-\infty}^{\infty}\frac{d}{dt}\hat{u}_j(t)e^{ij2\pi x} &= \left(\sum_{j=-\infty}^{\infty}\hat{u}_j(t)\hat{J}_j e^{ij2\pi x} - \sum_{j=-\infty}^{\infty}\hat{u}_j(t)e^{ij2\pi x}\right) \\ \frac{d}{dt}\hat{u}_j(t) &= (\hat{J}_j - \hat{J}_0)\hat{u}_j(t) = \hat{q}_j\hat{u}_j(t)\end{aligned}$$

with $\hat{q}_j = (\hat{J}_j - \hat{J}_0)$ where $\hat{J}_j$ is the jth Fourier coefficient of the kernel function $J(x)$. Solving the above equation we have

$$\hat{u}_j(t) = e^{\hat{q}_j t}\hat{u}_j(0),$$

where

$$\hat{u}_j(0) = \int_0^1 u(x,0)e^{-ij2\pi x}dx.$$

Thus the exact solution of the IDE (1) can be written as

$$u(x,t) = \sum_{j=-\infty}^{\infty} \hat{u}_j(0)e^{\hat{q}_j t}e^{ij2\pi x}. \tag{13}$$

Now taking the inverse of the discrete Fourier transform in (10) we get the approximate solution of (1) as

$$U_j^n = \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} g^n(h,\Delta t,k)\tilde{U}_k^0 e^{ik2\pi x_j}. \tag{14}$$

From (13) and (14)

$$\begin{aligned} u(x_j,t_m) - U_j^m &= \sum_{k=-\infty}^{\infty} \hat{u}_k(0)e^{\hat{q}_k t_m}e^{i2\pi k x_j} - \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} g^m(h,\Delta t,k)\tilde{U}_k^0 e^{ik2\pi x_j} \\ &= \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left(\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m(h,\Delta t,k)\tilde{U}_k^0\right) e^{i2\pi k x_j} \\ &\quad + \sum_{|k|>\frac{N}{2}} \hat{u}_k(0)e^{\hat{q}_k t_m}e^{ik2\pi x_j} \end{aligned} \tag{15}$$

Now taking the inner product on (15) and applying Parseval's relation

$$\begin{aligned} \|u(x_j,t_m) - U_j^m\|_h^2 &\le \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m(h,\Delta t,k)\tilde{U}_k^0\right|^2 \\ &\quad + \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m}\right|^2. \end{aligned} \tag{16}$$

The Poisson summation formula gives

$$\tilde{U}_k^m = \sum_{r=-\infty}^{\infty} \hat{u}_{k+rN}^m = \sum_{r=-\infty}^{\infty} \hat{u}_{k+rN}^{\infty,m}$$

where $\hat{u}_{k+rN}^m$ is the coefficient of the Fourier series at t_m for the periodic function u whereas $\hat{u}_{k+rN}^{\infty,m}$ represents the CFT for the nonperiodic (infinite-dimensional) case. A calculation similar to that leading to (9) gives

$$\hat{u}_k(0) = \int_0^1 u(x,0)e^{-ik2\pi x}dx = \hat{u}_k^0 = \hat{u}_0^\infty(2\pi k) = \hat{u}_k^{\infty,0}. \tag{17}$$

So the first part of the right-hand side of (16) can be written as

$$\begin{aligned}\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_j t_m} - g^m \tilde{U}_k^0\right|^2 &\leq \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|\left(e^{\hat{q}_j t_m} - g^m\right)\hat{u}_k^{\infty,0}\right|^2 \\ + \quad & \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|g^m(h,\Delta t,k)\sum_{r\neq 0}\hat{u}^{\infty,0}\left(2\pi(k+rN)\right)\right|^2 . \quad (18)\end{aligned}$$

Before giving the main convergence results let us introduce a lemma with reasonable restrictions on $J(x)$.

Lemma 3. *If $J(x)$ is a nonnegative even periodic function in $[0,1]$ and monotone-decreasing in $[0,\frac{1}{2}]$ then $\hat{q}_j \leq 0$ and $|\hat{J}_0 - \hat{J}_j| \leq \frac{1}{2}$ where $\hat{q}_j = (\hat{J}_j - \hat{J}_0)$ with $\hat{J}_j = \int_0^1 J(x)e^{-ij2\pi x}dx$.*

Proof. The proof of Lemma 3 follows directly from the definition of $\hat{J}_j$. ☐

Now

$$|e^{\hat{q}_j \Delta t}| \leq 1 \quad \text{and} \quad |g(h,\Delta t,k)| \leq 1.$$

So

$$|e^{\hat{q}_k t_m} - g^m(h,\Delta t,k)| = |\left(e^{\hat{q}_k \Delta t}\right)^m - g^m(h,\Delta t,k)| \leq m|e^{\hat{q}_k \Delta t} - g(h,\Delta t,k)|.$$

Also,

$$\begin{aligned} & e^{\hat{q}_k \Delta t} - g(h,\Delta t,k) \\ = \;& e^{\Delta t\left(\hat{J}_k - \hat{J}_0\right)} - \left(1 + \Delta t\left(\tilde{J}_k - \tilde{J}_0\right)\right) \\ = \;& \Delta t\left((\hat{J}_k - \hat{J}_0) - \left(\tilde{J}_k - \tilde{J}_0\right)\right) + \sum_{j=2}^{\infty}\frac{\Delta t^j}{j!}\left(\hat{J}_k - \hat{J}_0\right)^j , \qquad (19)\end{aligned}$$

and thus there exists $C_1(h)$ and C_2 (using (7) and (9)) such that

$$\left|e^{\hat{q}_j \Delta t} - g(h,\Delta t,k)\right| \leq \Delta t\left(C_1(h) + C_2\Delta t\right).$$

Here $C_1(h) \to 0$ as $h \to 0$, and C_2 is bounded, see [4] for exact detail. Thus we get the following bound.

Theorem 2. *Accuracy: If $J(x) \in H^r(1)$, $r > \frac{1}{2}$ satisfies **A1** - **A5** then there exist $C_1(h)$ and C_2 such that*

$$\left|e^{\hat{q}_j \Delta t} - g(h,\Delta t,k)\right| \leq \Delta t\left(C_1(h) + C_2\Delta t\right).$$

Applying Theorem 2 on the first part of the right-hand side of (18) we get

$$\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|(e^{\hat{q}_j t_m} - g^m)\hat{u}_0^{\infty}\right|^2 \leq \Delta t^2 \left(C_1(h) + C_2 \Delta t\right)^2 \|u_0\|^2. \tag{20}$$

Definition 1. *[1, page 223] For integer $k \geq 0$, $H^k(2\pi)$ is defined to be the closure of $C_p^k(2\pi)$ under the inner product norm*

$$\|\varphi\|_{H^k} = \left[\sum_{j=0}^{k} \|\varphi^{(j)}\|_{L^2}^2\right]^{\frac{1}{2}}.$$

For arbitrary real $s \geq 0$, $H^s(2\pi)$ can also be obtained following [1, pages, 219-223].

Theorem 3. *[1, page 223] For $s \in \mathbb{R}$, $H^s(2\pi)$ is the set of all series*

$$\varphi(x) = \sum_{m=-\infty}^{\infty} a_m \psi_m(x)$$

for which

$$\|\varphi\|_{*,s}^2 = |a_0|^2 + \sum_{|m|>0} |m|^{2s} |a_m|^2 < \infty$$

where

$$\psi_m(x) = \frac{1}{\sqrt{2\pi}} e^{imx}, \qquad m = 0, \pm 1, \pm 2, \cdots.$$

Moreover, the norm $||\varphi||_{*,s}$ is equivalent to the standard Sobolev norm $||\varphi||_{H^s}$ for $\varphi \in H^s(2\pi)$.

For exact details of the Theorem 3 please see [1, page 223]. In this section, to use norm definitions in a periodic domain $[0,1]$, we use $\|\varphi\|_{H^s} = \|\varphi\|_{H^s(1)}$.

Now

$$\begin{aligned}\left|\sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right| &\leq \sum_{s\neq 0} \left|\hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right| \\ \leq & \sqrt{\sum_{s\neq 0} \left|\hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\left(2\pi(k+sN)\right)^{\sigma}\right|^2} \sqrt{\sum_{s\neq 0} \left|\left(2\pi(k+sN)\right)\right|^{-2\sigma}}\end{aligned}$$

using the Cauchy-Schwartz inequality. Thus,

$$\begin{aligned}\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left| g^m(h,\Delta t,k) \sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2 &\leq \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left| \sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2 \\ \leq \quad \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} &\left(\sqrt{\sum_{s\neq 0} \left|\hat{u}^0\left(2\pi(k+sN)\right)\left(2\pi(k+sN)\right)^\sigma\right|^2}\right)\left(\sqrt{\sum_{s\neq 0} \left|\left(2\pi(k+sN)\right)\right|^{-2\sigma}}\right)\end{aligned}$$

gives

$$\begin{aligned}&\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left| g^m(h,\Delta t,k) \sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2 \\ \leq \quad & h^{2\sigma} C(\sigma) \sum_{|k|>\frac{N}{2}} |2\pi k|^{2\sigma} \left|\hat{u}^0\left(2\pi k\right)\right|^2 \\ \leq \quad & C(\sigma) h^{2\sigma} \|u_0\|^2_{*,\sigma} \equiv C(\sigma) h^{2\sigma} \|u_0\|^2_{H^\sigma(1)} \qquad (21)\end{aligned}$$

for some $\sigma > \frac{1}{2}$. Thus applying above bounds, Lemma 3 and Theorem 3 in (16) one gets

$$\begin{aligned}\|u(x_j,t_m) - U_j^m\|_h^2 \quad &\leq \quad \Delta t^2 \left(C_1(h) + C_2 \Delta t\right)^2 \|u_0\|^2 + C_3 h^{2\sigma} \|u_0\|^2_{H^\sigma(1)} + \sum_{|k|>\frac{N}{2}} |\hat{u}_k(0)|^2 \\ &\leq \quad \Delta t^2 \left(C_1(h) + C_2 \Delta t\right)^2 \|u_0\|^2 + C_3 h^{2\sigma} \|u_0\|^2_{H^\sigma(1)} \\ &\qquad + \left(\frac{2}{N}\right)^{2\sigma} \sum_{|k|>\frac{N}{2}} |k|^{2\sigma} \left|\hat{u}_0\left(2k\pi\right)\right|^2 \\ &\leq \quad \Delta t^2 \left(C_1(h) + C_2 \Delta t\right)^2 \|u_0\|^2 + C_3 h^{2\sigma} \|u_0\|^2_{H^\sigma(1)}, \end{aligned} \qquad (22)$$

for some $\sigma > \frac{1}{2}$. Thus we conclude this discussion with the following result.

Theorem 4. *If the approximation (6) of the initial value problem (1) is stable, $J, \hat{J}_k$, $-\frac{N}{2}+1 \leq k \leq \frac{N}{2}$ satisfy **A1** - **A5** and $u_0 \in H^\sigma(1)$ with $\sigma > \frac{1}{2}$, then there exist constants $C_1(h)$, C_2, $C_3(\sigma)$ such that*

$$\|u(x,t_m) - U_j^m(x)\| \leq \Delta t \left(C_1(h) + C_2 \Delta t\right) \|u_0\| + C_3(\sigma) h^\sigma \|u_0\|_{H^\sigma(1)}.$$

5 Convergence analysis for non-smooth initial data

Now let us consider an initial function which is not smooth enough so that $\|u_0\|_{H^{\nu}(1)}$ is bounded when $\nu > \frac{1}{2}$, but there exists $\nu_1 < \nu$ such that $\|u_0\|_{H^{\nu_1}(1)} < \infty$ and there exists α such that

$$\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|\hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2 \leq \Delta x^{2\alpha} C(u_0). \tag{23}$$

For example of such functions please see [4, 15] and references there in. We start varying (16) as

$$\begin{aligned}
\|u(x_j,t_m) - U_j^m\|_h^2 \;\leq\;& \sum_{|k|\leq M} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \tilde{U}_k^0\right|^2 \\
&+ \sum_{M\leq|k|\leq\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \tilde{U}_k^0\right|^2 + \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m}\right|^2
\end{aligned}$$

for some $0 < M < \frac{N}{2}$. Now with the same algebraic operations as we performed for the Theorem 4

$$\begin{aligned}
\|u(x_j,t_m) - U_j^m\|_h^2 \;\leq\;& \sum_{|k|\leq M} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \hat{u}_k^{\infty,0}\right|^2 + \sum_{M\leq|k|\leq\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \hat{u}_k^{\infty,0}\right|^2 \\
&+ \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m}\right|^2 + \sum_{|k|\leq M} \left|g^m \sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2 \\
&+ \sum_{M\leq|k|\leq\frac{N}{2}} \left|g^m \sum_{s\neq 0} \hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2,
\end{aligned}$$

$$\sum_{|k|\leq M} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \hat{u}_k^{\infty,0}\right|^2 \leq t^2(C_1(h) + C_2\Delta t)^2 \|u_0\|^2,$$

and

$$\begin{aligned}
& \sum_{M\leq|k|\leq\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t_m} - g^m \hat{u}_k^{\infty,0}\right|^2 + \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{q t_m}\right|^2 \\
=\;& C \sum_{|k|>M} \left|\hat{u}^{\infty,0}\right|^2 \leq C\left(\frac{1}{M}\right)^{2\nu_1} \sum_{|k|>M} |k|^{2\nu_1} \left|\hat{u}^{\infty,0}\right|^2 \leq C_3(\nu_1)\Delta x^{2\nu_1} \|u_0\|_{H^{\nu_1}(1)}
\end{aligned}$$

hold. Also following [4, 15], there exists $\alpha \in \mathbb{R}$ such that

$$\sum_{|k|\leq M}\left|g^m\sum_{s\neq 0}\hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2+\sum_{M\leq|k|\leq\frac{N}{2}}\left|g^m\sum_{s\neq 0}\hat{u}^{\infty,0}\left(2\pi(k+sN)\right)\right|^2$$

$$\leq \sum_{|k|\leq\frac{N}{2}-1}\left|\sum_{s\neq 0}\hat{u}^{\infty,0}\left(\frac{2\pi(k+sN)}{2L}\right)\right|^2\leq \Delta x^{2\alpha}C(u_0),$$

gives

$$\|u(x_j,t_m)-U_j^m\|^2\leq t^2(C_1(h)+C_2\Delta t)^2\|u_0\|^2+C_3(\nu_1)\Delta x^{2\nu_1}\|u_0\|_{H^{\nu_1}(1)}+\Delta x^{2\alpha}C(u_0).$$

Thus we finish this section with the following theorem.

Theorem 5. *If the initial value problem (1) with some non-smooth initial function* $u_0(x,0)=g(x)$ *is considered so that the condition (23) holds and is approximated by the stable one step finite difference formula (6),* $J, \hat{J}_k$, $-\frac{N}{2}+1\leq k\leq\frac{N}{2}$ *satisfy* ***A1*** *-* ***A5****, then*

$$\|u(x_j,t_m)-U_j^m\|\leq t(C_1(h)+C_2\Delta t)\|u_0\|+C_2(\nu_1)\Delta x^{\nu_1}\|u_0\|_{H^{\nu_1}(1)}+\Delta x^{\alpha}C_3(u_0).$$

where $C_1(h)$, C_2 *and* C_3 *are constants.*

6 Convergence of semidiscrete approximation

Applying the discrete Fourier transform to (5)

$$\frac{d}{dt}\tilde{U}_k=\tilde{q}_k\tilde{U}_k \tag{24}$$

where $\tilde{q}_k=\left(\tilde{J}_k-\tilde{J}_0\right)$ and its solution can be found as

$$\tilde{U}_k(t)=e^{\tilde{q}_kt}\tilde{U}_k(0). \tag{25}$$

Applying the inverse of the DFT in (25)

$$U(x_j,t)=\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}}e^{\tilde{q}_kt}\tilde{U}_k(0)e^{ik2\pi x_j}. \tag{26}$$

Thus comparing (13) and (26)

$$\begin{aligned}
u(x_j,t)-U(x_j,t) &= \sum_{k=-\infty}^{\infty} \hat{u}_k(0)e^{\hat{q}_k t}e^{ik2\pi x_j} - \sum_{k=-\frac{N}{2}-1}^{\frac{N}{2}} e^{\tilde{q}_k t}e^{ik2\pi x_j}\tilde{U}_k(0) \\
&= \sum_{k=-\frac{N}{2}-1}^{\frac{N}{2}} e^{ik2\pi x_j}\left(\hat{u}_k(0)e^{\hat{q}_k t} - e^{\tilde{q}_k t}\tilde{U}_k(0)\right) + \sum_{|k|>\frac{N}{2}} \hat{u}_k(0)e^{\hat{q}_k t}e^{ik2\pi x_j}.
\end{aligned} \tag{27}$$

Applying Parseval's relation to (27)

$$\begin{aligned}
\left|u(x_j,t)-U(x_j,t)\right|^2 &\le \sum_{k=-\frac{N}{2}-1}^{\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t} - e^{\tilde{q}_k t}\tilde{U}_k(0)\right|^2 + \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t}e^{ik2\pi x_j}\right|^2 \\
&\le \sum_{k=-\frac{N}{2}-1}^{\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t} - e^{\tilde{q}_k t}\tilde{U}_k(0)\right|^2 + \sum_{|k|>\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t}\right|^2 \qquad (28)
\end{aligned}$$

Now applying Poisson's formula to the first part of the right-hand side of (28),

$$\begin{aligned}
\sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|\hat{u}_k(0)e^{\hat{q}_k t} - e^{\tilde{q}_k t}\tilde{U}_k(0)\right|^2 &= \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \left|e^{\hat{q}_k t} - e^{\tilde{q}_k t}\right|^2 \left|\hat{u}_0^{\infty}(2\pi k)\right|^2 \\
&\quad + \sum_{k=-\frac{N}{2}+1}^{\frac{N}{2}} \sum_{s\neq 0} e^{2\tilde{q}_k t}\left|\hat{u}_0^{\infty}\left(2\pi(k+sN)\right)\right|^2.
\end{aligned}$$

Now

$$|e^{\hat{q}_k t} - e^{\tilde{q}_k t}|^2 = |e^{\hat{q}_k t}\left(1-e^{t(\tilde{q}_k-\hat{q}_k)}\right)|^2 \le t^2|\hat{q}_k - \tilde{q}_k|^2.$$

Now

$$\begin{aligned}
\tilde{q}_k - \hat{q}_k &= \left(\sum_{m=-\infty}^{\infty} \hat{q}_{k+mN}\right) - \hat{q}_k, \quad \text{using (7)} \\
&= \sum_{m\neq 0} \hat{q}_{k+mN} = \sum_{m\neq 0}\left(\hat{J}_{k+mN} - \hat{J}_{mN}\right), \quad \text{since } \hat{q}_k = \hat{J}_k - \hat{J}_0, \\
&= \sum_{m\neq 0}\left(\hat{J}^{\infty}(2\pi(k+mN)) - \hat{J}^{\infty}(2\pi mN)\right), \quad \text{using (9).}
\end{aligned}$$

So from [4] it follows that

$$|\tilde{q}_k - \hat{q}_k| \le 2\hat{J}^\infty\left(\frac{\pi}{h}\right) = C_1(h),$$

and $C_1(h) \to 0$ as $h \to 0$ when J^∞ is smooth enough (if $J^\infty \in L_2(\mathbb{R})$, then $|\hat{J}^\infty(\xi)| \to 0$ as $|\xi| \to \infty$ [14], [16, page 30]). Thus

$$|e^{\hat{q}_k t} - e^{\tilde{q}_k t}|^2 \le t^2 C_1^2(h).$$

So applying the relation (17), Parseval's relation and Theorem 3

$$\begin{aligned}\sum_{|k|\le\frac{N}{2}} \left|\left(e^{\hat{q}_k t} - e^{\tilde{q}_k t}\right)\hat{u}_0^\infty\right|^2 &\le t^2 C_1^2(h) \sum_{|k|\le\frac{N}{2}} |\hat{u}_k(0)|^2 \\ \le\ & t^2 C_1^2(h)\left(|\hat{u}_0(0)|^2 + \sum_{|k|>0} |k|^0\, |\hat{u}_k(0)|^2\right) \\ \le\ & t^2 C_1^2(h)\|u_0\|_{*,0}^2 \equiv t^2 C_1^2(h)\|u_0\|_{H^0(1)}^2 = t^2 C_1^2(h)\|u_0\|^2.\end{aligned}$$

Thus, similar to Section 4 there exists a constant $C_2(\sigma)$ such that

$$\|u(x_j,t) - U(x_j,t)\|^2 \le t^2 C_1^2(h)\|u_0\|^2 + C_2^2(\sigma) h^{2\sigma}\|u_0\|_{H^\sigma(1)}^2.$$

Thus we conclude

Theorem 6. *If the semidiscrete approximation (5) of the initial value problem (1) is stable* $J, \hat{J}_k$, $-\frac{N}{2}+1 \le k \le \frac{N}{2}$ *satisfy **A1 - A5** and* $u_0 \in H^\sigma(1)$ *with* $\sigma > \frac{1}{2}$, *then there exist constants* $C_1(h)$, $C_2(\sigma)$ *such that*

$$\|u(x_j,t) - U(x_j,t)\| \le tC_1(h)\|u_0\| + C_2(\sigma)h^\sigma\|u_0\|_{H^\sigma(1)}.$$

7 Numerical Experiments and discussion

Here we start by experimenting numerical error in the approximation (6) of (1). We compute numerical error at $t = 1$. Figure 2 shows the behaviour of $\|u(\cdot,t_n) - U^n(\cdot)\|_h$ for various choices of h and Δt with smooth and non-smooth u_0 for all $x \in [0,1]$ where $J^\infty(x) = \sqrt{\frac{10}{\pi}} exp(-10x^2)$. From Figure 2, we observe that for smooth u_0 the rate of convergence of the solutions are faster than for the non-smooth u_0. Here we also notice that choices of h and Δt have an impact on the rate of convergence. That



Figure 2: Here we plot the error term $\|u(\cdot,t_n)-U^n(\cdot)\|_h$ at $t=1$ by varying Δt and h with $J^\infty(x)=\sqrt{\frac{10}{\pi}}e^{-10x^2}$ and (a) $u_0(x)=\sqrt{\frac{1}{\pi}}e^{-(x-\frac{1}{2})^2}$ (left figure), (b) $u_0(x)=\frac{1}{2}e^{-|x-\frac{1}{2}|}$ (right figure).

agrees with our theoretical estimates (and also motivates us to investigate further the theoretical stability and convergence analysis of such a one step approximation). From this study we notice that the full discrete scheme is conditionally stable. The accuracy of the scheme depends on the smoothness of the initial function. We have some limitations in this study. We impose some reasonable restrictions on the kernel function to prove the stability and convergence results. The analysis of higher order schemes in one and multi-dimensions leaves as a future study which is of course more challenging.

References

[1] Kendall Atkinson and Weimin Han. *Theoretical Numerical Analysis*. Springer, 2001.

[2] Peter W. Bates and Adam Chmaj. A discrete convolution model for phase transitions. *Archive for Rational Mechanics and Analysis*, 150(4):281–305, 1999.

[3] Peter W. Bates, Paul C. Fife, Xiaofeng Ren, and Xuefeng Wang. Travelling waves in a convolution model for phase transitions. *Archive for Rational Mechanics and Analysis*, 138(2):105–136, July 1997.

[4] Samir K. Bhowmik. *Numerical approximation of a nonlinear partial integro-differential equation*. PhD thesis, Heriot-Watt University, Edinburgh, UK, April, 2008.

[5] Samir Kumar Bhowmik. Stability and convergence analysis of a one step approximation of a linear partial integro-differential equation. *Technical Report, KdV Institute for Mathematics, University of Amsterdam*, December 2009.

[6] C-K Chen and Paul C. Fife. Nonlocal models of phase transitions in solids. *Advances in Matheamtical Sciences and Applications*, 10:821–849, 2000.

[7] Fengxin Chen. Uniform stability of multidimensional travelling waves for the nonlocal Allen-Cahn equation. *Fifth Mississippi State Conference on Differential Equations and Computational Simulations, Electronic Journal of Differential Equations*, Conference 10:109–113, 2003.

[8] Daniel J. Duffy. *Finite Difference Methods for Financial Engineering: a Partial Differential Equation Approach*. Wiley Finance, 2006.

[9] Dugald B. Duncan, M. Grinfeld, and I. Stoleriu. Approximating a convolution model of phase separation. Conference Talk, Dundee, at May 21, 2003.

[10] Dugald B. Duncan, M Grinfeld, and I Stoleriu. Coarsening in an integro-differential model of phase transitions. *Euro. Journal of Applied Mathematics*, 11:511–523, 2000.

[11] Paul C. Fife. Models of phase separation and their Mathenmatics. *Electronic Journal of Differential Equation*, 48:1–26, 2000.

[12] Paul C. Fife. Well-posedness issues for medels of phase transitions with weak interaction. *Nonlinearity*, 14:221–238, 2001.

[13] Carlo R. Laing and William C. Troy. PDE methods for nonlocal models. *SIAM J. Applied dynamical systems*, 2(3):487–516, 2003.

[14] K. Maleknejad. A comparison of fourier extrapolation methods for numerical solution of deconvolution. *New York Journal of Mathematics*, 183:533–538, 2006.

[15] J. C. Strikwerda. *Finite Difference Schemes and Partial Differential Equations*. Wadsworth and Brooks, Cole Advanced Books and Software, Pacific Grove, California, 1989.

[16] L. N. Trefethen. *Spectral Methods in* MATLAB. SIAM, Philadelphia, 2000.